Journal of Immunological Methods, 138 (1991) 311–312

© 1991 Elsevier Science Publishers B.V. 0022-1759/91/$03.50

Contents to volume 138 (1991)

Number 1 – 8 April 1991

Research reports

Simplified quantitation of cytotoxicity by integration of specific lysis against effector cell concentration at a constant target cell concentration and measuring the area under the curve
J.F. Dye, S.S. Somers and P.J. Guillou (U.K.) 1

Proteolytic digestion of mouse IgE
S. Haba and A. Nisonoff (U.S.A.) 15

An avidin-biotin-peroxidase assay to detect synthetic peptides bound to polystyrene plates
G.E. Griesmann, D.J. McCormick and V.A. Lennon (U.S.A.) 25

Quantitation of biologically active IL-1 by a sensitive assay based on immobilized human IL-1 receptor type II ($IL\text{-}1R_{II}$)
G. Scapigliati, P. Bossu, S. Censini, A. Tagliabue, D. Boraschi and P. Ghiara (Italy) 31

A new radioimmunoassay for human mast cell tryptase using monoclonal antibodies
I. Enander, P. Matsson, J. Nystrand, A.-S. Andersson, E. Eklund, T.R. Bradford and L.B. Schwartz (Sweden, U.S.A.) 39

Sensitive ELISA for interleukin-6. Detection of IL-6 in biological fluids: synovial fluids and sera
M. Helle, L. Boeije, E. de Groot, A. de Vos and L. Aarden (The Netherlands) 47

Assay restriction profiles of three monoclonal antibodies recognizing the G3m(u) allotype. Development of an allotype specific assay
P.N. Nelson, S.M. Fletcher, D. MacDonald, D.M. Goodall and R. Jefferis (U.K.) 57

Description of a urease-based microELISA for the analysis of limiting dilution microcultures
M.C. Cerrone and R.E. Kuhn (U.S.A.) 65

A method for generating antigen-specific rat T helper cell clones
J. Katz and S.M. Michalek (U.S.A.) 77

Determination of urinary Tamm-Horsfall protein by ELISA using a maleimide method for enzyme-antibody conjugation
I. Uto, T. Ishimatsu, H. Hirayama, S. Ueda, J. Tsuruta and T. Kambara (Japan) 87

Direct time-resolved fluorescence immunoassay for serum oestradiol based on the idiotypic anti-idiotypic approach
A. Altamirano-Bustamante, G. Barnard and F. Kohen (Israel, U.K.) 95

Analysis of lymphocyte shape by visual classification, calculated measures of shape or light scattering
S. Eisele, J.M. Lackie, H. Riedwyl, A. Zimmermann and H.U. Keller (Switzerland, U.K.,) 103

Rapid whole blood assay for HIV-1 seropositivity using an Fab-peptide conjugate
K.M. Wilson, M. Gerometta, D.B. Rylatt, P.G. Bundesen, D.A. McPhee, C.J. Hillyard and B.E. Kemp (Australia) 111

Short communications

Double wavelength measurement of 3,3′,5,5′-tetramethylbenzidine (TMB) provides a three-fold enhancement of the ELISA measuring range
S. Madersbacher and P. Berger (Austria) 121

A cyclized peptide for studies of human parvovirus B19 infection
E. Fridell, J. Trojnar, H. Mehlin and B. Wahren (Sweden) 125

Letters to the editors

Chemiluminescent quantitation of lymphocyte surface antigens
E. Roffman and N. Frenkel (U.S.A.) 129

Flow cytometric measurement of the respiratory burst activity of phagocytes using dihydrorhodamine 123
G. Rothe, A. Emmendörffer, A. Oser, J. Roesler and G. Valet (F.R.G.) 133

Corrigendum

Methodological considerations when using nitrocellulose immunoblotting from polyacrylamide gels to study the mould allergens *Aspergillus fumigatus* and *Alternaria alternata*
Å. Karlsson-Borgå and W. Rolfsen (Sweden) 137

Announcements 139

Notes to authors 141

Number 2 – 25 April 1991

Research reports

Improved permeabilization procedure for flow cytometric detection of internal antigens. Analysis of interleukin-2 production
M. Labalette-Houache, G. Torpier, A. Capron and J.P. Dessaint (France) 143

Computerised transfer and processing of data from experiments measuring cellular proliferation by incorporation of tritiated thymidine
K. Schaudt and G. Pawelec (F.R.G.) 155

A method to differentiate between anti-C1q antibodies and C1q-binding immune complexes using collagenase-digested solid phase C1q
J.E. Menzel, O. Scherak, G. Kolarz, F. Gamerith and U. Youngchaiyud (Austria, Thailand) 165

Isolation of IgG1-secreting switch variants from IgM hybridomas produced after in vitro immunization
M. Mercken, M. Vandermeeren, U. Lubke, J. Boons, A. Van de Voorde and J. Gheuens (Belgium) 173

Purification and stabilisation of a poorly soluble mouse IgG3 monoclonal antibody
W. Jiskoot, A.-M.V. Hoven, A.A.M. De Koning, M.F. Leerling, C.H.K. Reubsaet, D.J.A. Crommelin and E.C. Beuvery (The Netherlands) 181

Fetal intestinal transplant model for mucosal immune responses
A.G. Kantak and M.S. Cooperstock (U.S.A.) 191

Detection of rheumatoid-like factors in serum of chickens immunized with bacterial immunoglobulin binding proteins
R. Raeder and M.D.P. Boyle (U.S.A.) 201

Production of a hybridoma cell line secreting retinoic acid-specific monoclonal antibody
H.-R. Zhou, M.M. Abouzied and M.H. Zile (U.S.A.) 211

ELISA for the detection of anticardiolipin antibodies. High specificity based on the use of adult bovine serum as buffer and systematic subtraction of non-specific binding
A. Rupin, Y. Gruel, H. Watier, A.C. Girard, J. Leroy and P. Bardos (France) 225

Preparation of europium-streptavidin in a time-resolved fluoroimmunoassay for interleukin-3
H.-P. Knopf and R. Papoian (Switzerland) 233

The use of adenovirus-infected HeLa cells for the detection of low titer autoantibodies
R. Slobbe, B. Van Esch, T. Kveder and W.J. Van Venrooij (The Netherlands, Yugoslavia) 237

The influence of epitope density on the estimation of the affinity of antibody for complex antigens
G.P. Holland and M.W. Steward (U.K.) 245

Simultaneous identification of eight leucocyte subsets of human peripheral blood using three-colour immunofluorescence flow cytometric analysis
P. Carayon, A. Bord and M. Raymond (France) 257

Purification of human complement factor D from the peritoneal fluid of patients on chronic ambulatory peritoneal dialysis
E. Catana and J.A. Schifferli (Switzerland) 265

Preparation of clinical grade monoclonal antibodies from serum-containing cell culture supernatants
W. Jiskoot, J.J.C.C. Van Hertrooij, A.-M. V. Hoven, J.W.T.M. Klein Gebbinck, T. Van der Velden-de Groot, D.J.A. Crommelin and E.C. Beuvery (The Netherlands) 273

Detection of eosinophil cationic protein (ECP) by an enzyme-linked immunosorbent assay
C.M. Reimert, P. Venge, A. Kharazmi and K. Bendtzen (Denmark, Sweden) 285

Optimum dissociating condition for immunoaffinity and preferential isolation of antibodies with high specific activity
V.C.W. Tsang and P.P. Wilkins (U.S.A.) 291

Letter to the editors

A rapid colorimetric bioassay for transforming growth factor β (TGF-β) using a microwell plate reader
M. Absher, L. Baldor and J. Kelley (U.S.A.) 301

Software notice

Automatic collection, transfer, and processing of cellular proliferation data
K. Schaudt and G. Pawelec (F.R.G.) 305

Book reviews 307

Announcement 309

Contents to volume 138 (1991) 311

Notes to authors 313